

SUBMILLIMETER GALAXY NUMBER COUNTS AND MAGNIFICATION BY GALAXY CLUSTERS

MARCOS LIMA, BHUVNESH JAIN, MARK DEVLIN, AND JAMES AGUIRRE
Department of Physics & Astronomy, University of Pennsylvania, Philadelphia PA 19104
Draft version April 28, 2010

ABSTRACT

We present an analytical model which reproduces measured galaxy number counts from surveys in the wavelength range of 500 μm to 2 mm. The model involves a single high-redshift galaxy population with a Schechter luminosity function which has been gravitationally lensed by galaxy clusters in the mass range 10^{13} to $10^{15} M_{\odot}$. This simple model reproduces both the low flux and the high flux end of the number counts reported by the BLAST, SCUBA, AzTEC and the SPT surveys. In particular, our model accounts for the most luminous galaxies detected by SPT as the result of high magnifications by galaxy clusters (magnification factors of 10-30). This interpretation implies that submillimeter and millimeter surveys of this population may prove to be a useful addition to ongoing cluster detection surveys. The model also implies that the bulk of submillimeter galaxies detected at wavelengths larger than 500 μm lie at redshifts greater than 2.

Subject headings: galaxies: clusters — gravitational lensing — submillimeter: galaxies

1. INTRODUCTION

Over the last decade, submillimeter (submm) surveys have yielded significant advances in our understanding of the galaxy population responsible for the high-redshift component of the cosmic infrared background (CIB) (Fixsen et al. 1996; Smail et al. 1997; Hughes et al. 1998; Barger et al. 1998; Dwek et al. 1998; Fixsen et al. 1998; Greve et al. 2004; Pope et al. 2006; Coppin et al. 2006; Devlin et al. 2009). With typical far-infrared (FIR) luminosities $> 10^{12} L_{\odot}$, submm galaxies are presumed to be the high-redshift counterparts to (ultra) luminous infrared galaxies (LIRGs, ULIRGs). The high luminosity of these galaxies is the result of star formation rates of 100–1000 $M_{\odot} \text{yr}^{-1}$. Approximately half of these galaxies are located at $1.9 \lesssim z \lesssim 2.9$ (Chapman et al. 2005; Aretxaga et al. 2007), dominating the total star formation rate at this epoch (Pérez-González et al. 2005).

One way to express the results of submm surveys is through number counts of galaxies as a function of flux for each observed wavelength. The shape of these counts has been interpreted as arising from different populations of galaxies whose characteristics evolve over cosmic time (Lagache et al. 2003, 2004; Pearson & Khan 2009; Le Borgne et al. 2009). These empirical models have successfully reproduced the counts. However, they may be masking a simpler explanation for the departure of the counts from a Schechter distribution at the high-flux end: magnification due to high redshift galaxy clusters and groups (e.g. Blain 1996; Perrotta et al. 2002; Negrello et al. 2007).

Millimeter wavelength surveys have also aimed at detecting galaxy clusters via the Sunyaev Zel’dovich (SZ) effect (Hincks et al. 2008; Carlstrom et al. 2009); the first results, including CMB power spectra and cluster catalogs, have been released recently (Fowler et al. 2010; Staniszewski et al. 2009; Vanderlinde et al. 2010). The number of detected clusters remains relatively low, primarily due to the low value of σ_8 . However, other ef-

fects could be reducing the sensitivity of the surveys (e.g. Lima et al. 2009, 2010).

The South Pole Telescope (SPT) has measured number counts of dusty galaxies at wavelengths $\lambda = 1.4$ mm and 2.0 mm over an area of 87 deg² (Vieira et al. 2009). The observed numbers at the bright end are higher than expected: these galaxies are either at high redshifts and intrinsically exceptionally luminous, or have been magnified by gravitational lensing, or are simply at much lower redshifts than the bulk of the population of submm galaxies. The latter possibility is disfavored by the lack of detected counterparts in other surveys that probed the low redshift population (Vieira et al. 2009). The possibility that these galaxies are at high redshifts and intrinsically bright would require them to be far more luminous than an underlying Schechter-like luminosity function would permit. Thus the favored explanation is that they have typical luminosities for high- z galaxies, but have been magnified by foreground galaxies or clusters. Lensing of high-redshift background submm galaxies has been observed (Smail et al. 1997, 2002), and more recently in the Bullet Cluster with the 1.1 mm AzTEC receiver on the Atacama Submillimeter Telescope Experiment as well as with the Balloon-borne Large Aperture Submillimeter Telescope - BLAST (Wilson et al. 2008; Rex et al. 2009; Gonzalez et al. 2009).

In this *Letter*, we explore the possibility that the existing observed galaxy number counts over a wide range of wavelengths can be reproduced by a *single* population of galaxies at high-redshift. Foreground galaxy groups and clusters gravitationally lenses the background submm population (Lima et al. 2009) and leads to significant enhancements of the high-flux end to the galaxy counts. In § 2 we describe the lensing magnification formalism, which we then apply to a high- z galaxy population and present results in § 3. We discuss implications for high- z galaxies and the cluster searches in § 4.

Throughout, we use a fiducial cosmology for a flat universe with parameter values based on the results of the Wilkinson Microwave Anisotropy Probe fifth year data




Figure 1. Intrinsic and lensed number counts dn/dS for a Schechter function describing galaxies at different redshifts. Also shown are the observed counts for SPT *dusty* submm galaxies at $\lambda = 2.0$ mm, after removal of low-redshift galaxies with IRAS counterparts.

release (WMAP5, Komatsu et al. 2009). The cosmological parameters (and their values) are the normalization of the initial curvature spectrum $\delta_\zeta (= 2.41 \times 10^{-4})$ at $k = 0.02 \text{ Mpc}^{-1}$ (corresponding to $\sigma_8 = 0.8$), its tilt $n (= 0.96)$, the baryon density relative to critical $\Omega_b h^2 (= 0.023)$, the matter density $\Omega_m h^2 (= 0.13)$, and two dark energy parameters: its density $\Omega_{\text{DE}} (= 0.74)$ and equation of state $w (= -1)$, which we assume to be constant. Since lensing effects depend on the abundance of dark matter halos, which is exponentially sensitive to linear matter perturbations, we also consider changes in σ_8 consistent with the WMAP5 errors of $\Delta\sigma_8 \approx 0.03$. Our central value and uncertainty for σ_8 is consistent with the WMAP7 analysis (Komatsu et al. 2010).

2. NUMBER COUNTS WITH LENSING MAGNIFICATION

In recent papers (Lima et al. 2009; Jain & Lima 2010) we have presented a halo model for calculating the effects of lensing magnification by galaxy groups and clusters. Here we specialize to the case of steep galaxy counts at high redshifts, where lensing effects are quite dramatic. We assume a Schechter function (Schechter 1976) for the intrinsic number density distribution of a population of galaxies

$$\frac{dn}{dS} = \frac{n^*}{S^*} \left(\frac{S}{S^*}\right)^\alpha e^{-S/S^*}, \quad (1)$$

where n^* , S^* and α are free parameters. Due to lensing magnification by intervening halos, the intrinsic dn/dS is changed to its observed counterpart as

$$\frac{dn_{\text{obs}}(S_{\text{obs}})}{dS_{\text{obs}}} = \int d\mu \frac{P(\mu)}{\mu} \frac{dn}{dS} \left(\frac{S_{\text{obs}}}{\mu}\right), \quad (2)$$

where μ is the lensing magnification and $P(\mu)$ is its probability for a given galaxy population at redshift z_s . Conditional probabilities quantify the effects of different magnification ranges on the observed flux density S_{obs} . The integrand of Eq. 2 defines the probability $P(\mu|S_{\text{obs}})$

$$P(\mu|S_{\text{obs}}) = \left(\frac{dn_{\text{obs}}(S_{\text{obs}})}{dS_{\text{obs}}}\right)^{-1} \frac{P(\mu)}{\mu} \frac{dn}{dS} \left(\frac{S_{\text{obs}}}{\mu}\right), \quad (3)$$

which can be interpreted as the relative contribution of a given μ to the total $dn_{\text{obs}}/dS_{\text{obs}}$ at S_{obs} (Paciga et al. 2008). Similarly $P(\mu_{\text{min}}|S_{\text{obs}}) = \int_{\mu_{\text{min}}}^{\infty} d\mu P(\mu|S_{\text{obs}})$ measures the integrated contribution from all $\mu > \mu_{\text{min}}$. The mean magnification at a given S_{obs} is defined as

$$\langle\mu\rangle(S_{\text{obs}}) = \int_0^{\infty} d\mu \mu P(\mu|S_{\text{obs}}). \quad (4)$$

The distribution $P(\mu)$ can be estimated either by ray-tracing on N-body simulations (e.g. Hilbert et al. 2007), or by semi-analytical methods (e.g. Perrotta et al. 2002; Lima et al. 2009), integrating halo contributions on the line of sight up to the source redshift

$$P(>\mu) = \int_0^{z_s} dz_l \frac{D_A^2(z_l)}{H(z_l)} \int_{M_{\text{th}}}^{\infty} d \ln M \frac{dn(z_l, M)}{d \ln M} \Delta\Omega_\mu, \quad (5)$$

where D_A is the angular diameter distance, H is the Hubble parameter, $dn/d \ln M$ is the halo mass-function and $\Delta\Omega_\mu = \Delta\Omega_\mu(z_s, z_l, M)$ is the cross-section for magnifications larger than μ produced by halos of mass M at redshift z_l on sources at redshift z_s . The integrand

$$\frac{d^2 P(>\mu)}{d \ln M dz_l} = \frac{D_A^2(z_l)}{H(z_l)} \frac{dn(z_l, M)}{d \ln M} \Delta\Omega_\mu, \quad (6)$$

gives the range of halo masses and redshifts contributing most to the probability of a specified minimum magnification μ .

In summary, Eqs. 2 and 5 give the total effect on the counts, Eq. 3 indicates which magnifications contribute most to the given S_{obs} and Eq. 6 tells us which halo masses and redshifts contribute to a given magnification.

We use this halo-model $P(\mu)$ and correct it for a number of effects. First, as described in Lima et al. (2009), we match our $P(\mu)$ at large magnifications to that of ray-tracing in dark matter simulations (Hilbert et al. 2007) by tuning the ellipticity of our halos. Next, we account for the effect of luminous matter, which can lead to higher densities via gas cooling, using the simulation results of Hilbert et al. (2008). We also correct for the combination of finite source size and multiple image effects. Magnification effects, especially for large magnifications, are sensitive to the value of σ_8 since it affects the abundance of cluster halos. In the next section we discuss how we account for the uncertainties in our model by giving a range for our predictions.

Our analytical calculation of $P(\mu)$ has some advantages over the approach of numerical simulations (we can easily study changes in source redshift, σ_8 and the contribution from different halo masses and redshift), but it also has some limitations. We only use the one-halo term, which is accurate at the high magnifications relevant for the effects considered here but overestimates the lensing




Figure 2. (*Left*) : Intrinsic and lensed dn/dS for a Schechter function describing galaxies at $z_s = 3.0$ and different wavelengths. Dashed lines indicate the intrinsic Schechter functions (with different S^* values) and the shaded regions display the range of lensing predictions, as described in the text. Also shown are observed counts for BLAST at $\lambda = 500 \mu\text{m}$, SCUBA at $\lambda = 850 \mu\text{m}$ and SPT at both $\lambda = 1.4 \text{ mm}$ and 2.0 mm . The SPT counts are for *dusty* galaxies, after removal of galaxies with synchrotron emission and galaxies with IRAS counterparts. Note that no similar removal has been applied to the BLAST data, which includes both high and low redshift galaxies. (*Right*) : Unified scaled curves showing $d\tilde{n}/d\tilde{S}$ and the various data points. The only parameter used in the scaling is S^* ; its values at the different wavelengths are given in Table 1 and are consistent with theoretical expectations.

Survey	$\lambda(\mu\text{m})$	$S^*(\text{mJy})$
BLAST	500	11.00
SCUBA	850	3.26
AzTEC	1100	1.28
SPT	1400	1.00
SPT	2000	0.35

Table 1

Scale S^* in the Schechter function for different surveys.

contribution at $\mu \sim 1$. We do not include a distribution of ellipticities or halo substructure, which can also increase magnification cross-sections. We have instead tuned the average halo ellipticity to match the $P(\mu)$ measured in dark matter simulations of Hilbert et al. (2007) (see Lima et al. (2009) for a detailed discussion). And whereas we account for the effects of baryons observed in numerical simulations by Hilbert et al. (2008), these authors note that their simulation still underestimate baryonic effects for halos of smaller masses as they do not predict sufficient numbers of multiply imaged quasars. Finally, the effect of finite source size is very uncertain given the lack of our knowledge about submm galaxies and its sensitivity to the precise caustic structure of the lenses. A proper inclusion of all these missing effects would likely increase the magnification probabilities compared to our current model.

3. RESULTS

In Fig. 1 we illustrate the lensing effect on an intrinsic Schechter function distribution given by Eq. 1 for sources at different redshifts. In all results presented here, we fix $\alpha = -1.0$ and $n^* = 5 \times 10^3 \text{ deg}^{-2}$. Changing to $\alpha = -1.5$ does not have a significant effect – while it

matches the faint end behavior of dn/dS for the model of Lagache et al. (2004), we preferred to use $\alpha = -1.0$ since this fits better the number counts at shorter wavelengths. With these parameter values, our counts are lower than the model of Lagache et al. (2004) at all S , which also ensures that the total flux does not exceed the CIB (Dwek et al. 1998; Fixsen et al. 1998).

Fig. 2 illustrates our main results: we show predicted number counts that include lensing (gray bands), assuming galaxies at $z_s = 3.0$, along with measured number counts from submm surveys at different wavelengths. As indicated in the panels, these are: BLAST at $\lambda = 500 \mu\text{m}$ (Devlin et al. 2009), SCUBA at $\lambda = 850 \mu\text{m}$ (Coppin et al. 2006), AzTEC at 1.1 mm (Austermann et al. 2010) and SPT *dusty* submm galaxies at $\lambda = 1.4 \text{ mm}$ and 2.0 mm (Vieira et al. 2009). In all our results, the SPT number counts correspond to those of Vieira et al. (2009), after *removing* both synchrotron emission galaxies as well as low-redshift galaxies that have matches with galaxies in the Infrared Astronomy Satellite survey (IRAS, Moshir et al. 1992; Fisher et al. 1995; Oliver et al. 1996). Predictions are shown by bands rather than curves to reflect the uncertainties in the model as discussed below. The intrinsic luminosity function is the Schechter function described above. All the data sets can be fit by changing the single parameter S^* once lensing magnification is included. This remarkable result implies that, within the measurement and theoretical uncertainties, a single high- z population of galaxies is sufficient to describe all the observations. The high flux measurements of BLAST and SPT are fit by highly magnified galaxies – if these counts were dominated by a population of a different galaxy type, it would be a co-




Figure 3. Probability $P(\mu_{\min}|S_{\text{obs}})$ of the minimum magnification μ_{\min} , given an observed flux density S_{obs} (left panel), and the corresponding average magnification $\langle\mu\rangle$ (right panel). We assume a Schechter function describing galaxies at $z_s = 3.0$ which, after lensing, predicts counts consistent with those of SPT dusty submm galaxies at $\lambda = 1.4$ mm (see Fig. 2).

incidence that their relative counts fit the same scaling with wavelength as the fainter (“normal”) population.

The lower bound for the predictions in Fig. 2 uses $\sigma_8 = 0.77$ while the upper bound uses $\sigma_8 = 0.83$ – these reflect the uncertainties in the WMAP results as discussed above. The enhancement due to baryons is factored into these predictions following Hilbert et al. (2008), though it is likely to be an underestimate of baryonic effects as discussed above in Section 2. The effect of source sizes and multiple imaging is uncertain; we simply assume that due to the finite source size the effective magnification is reduced by 50% to 25% (for the lower and upper bounds respectively)

The right panel of Fig. 2 shows all points rescaled by plotting

$$\frac{d\tilde{n}}{d\tilde{S}} = \tilde{S}^\alpha e^{-\tilde{S}}, \quad (7)$$

where $\tilde{n} = n/n^*$ and $\tilde{S} = S/S^*$. In Table 1, we show values of S^* used for each wavelength. We compared the frequency scaling of S^* with that of a typical Spectral Energy Distribution (SED) of submm galaxies $\text{SED}(\lambda) \propto \epsilon(\lambda)B(T, \lambda)$, with emissivity $\epsilon(\lambda) = 1 - \exp[-(\lambda_0/\lambda)^\beta]$ and blackbody spectrum $B(T, \lambda)$ at temperature T . The values of S^* are consistent with $\beta = 1 - 2$ and $T = 20 - 40$ K, as expected for high redshift submm galaxies.

The fit to the BLAST counts at $\lambda = 500, 350$ and $250 \mu\text{m}$ falls further below the high flux measurement at shorter wavelengths, suggesting the need for a lower redshift population. Indeed, Eales et al. (2009) have identified the radio and $24 \mu\text{m}$ counterparts of the bright BLAST sources. At $250 \mu\text{m}$ almost all of the bright sources are identified as being at redshifts lower than 1. One third to one half of the sources in the highest flux bin at $500 \mu\text{m}$ come from sources with $z < 1$. Removing these would lower the corresponding point in Fig. 2,

in agreement with the model. The remaining sources at $500 \mu\text{m}$ (and less than a tenth of the sources at $250 \mu\text{m}$) are likely the result of lensing. Similar results should be expected with the upcoming release of the large-area Herschel surveys.

Since the lensed distributions at $z_s = 3.0$ are consistent with SPT data points at both wavelengths, we study the range of magnifications and halo masses that would be contributing most in this case. As we consider larger values of S_{obs} , in particular $S_{\text{obs}}/S^* \gtrsim 10$, the observed sources come from intrinsically low flux sources which have been magnified significantly. Fig. 3 shows $P(\mu_{\min}|S_{\text{obs}})$ and indicates magnifications that contribute most at each S_{obs} . For instance, for $S_{\text{obs}} = 20 - 40$ mJy, the right panel shows that $\langle\mu\rangle \sim 20 - 30$. Another way to see this is via $P(\mu_{\min}|S_{\text{obs}})$ (left panel), which integrates out the effects above a certain μ_{\min} , and shows that for $S_{\text{obs}} = 20 - 40$ mJy, we have $P(\mu_{\min}|S_{\text{obs}}) > 0.5$ for $\mu_{\min} \sim 10 - 20$.

These results imply that magnifications of $10 - 30$ are necessary to explain the boost in dn/dS at $S_{\text{obs}} \sim 10 - 40$ mJy, if it is due to lensing of an intrinsic Schechter distribution. Note that due to the finite size of submm galaxies, their magnifications must have a cut-off, which is difficult to estimate, but may be in the range $\mu \sim 10 - 40$ (Perrotta et al. 2002). However, there is still substantial uncertainty in these estimates, since galaxies have been measured with estimated magnifications of at least ~ 45 (Paciga et al. 2008; Kneib et al. 2004).

In Fig. 4 we show $d^2P/d\ln M dz_l$ as a function of halo mass for different values of μ_{\min} and z_l . This indicates that halo masses above $10^{13}h^{-1}M_\odot$ contribute significantly to magnifications of $10 - 30$, with most of the contribution coming from $\sim 10^{14}h^{-1}M_\odot$. Note that this does not include baryonic effects, which boost the con-




Figure 4. The integrand $d^2P(>\mu)/d\ln M dz_l$ as a function of halo mass M for different values of μ and z_l . We have arbitrarily normalized these curves for better visualization. Note that the peak lensing contribution for source galaxies at $z_s = 3$ comes from lens halos at $z_l \sim 0.5$. This does not include baryonic effects, which boost the contribution from $M \lesssim 10^{13} h^{-1} M_\odot$ halos.

tribution from lower mass ($\lesssim 10^{13} h^{-1} M_\odot$) halos as discussed in Section 2.

We have assumed that all source galaxies are at a fixed redshift z_s , whereas in reality they have a redshift distribution which needs to be incorporated in the computation. A galaxy population at fixed redshift has number counts dn/dS that preserve the functional form of its luminosity function. Therefore our assumption of a Schechter form for the flux counts is consistent with a corresponding luminosity function at that redshift. The results at fixed z_s therefore can only provide lower and upper limits for the magnification effects. Nevertheless, our results imply that the bulk of the population of submillimeter galaxies are at redshifts $z \gtrsim 2$. A distribution of the galaxies over a range of redshift would not be able to produce the high flux measurements without a significant fraction at very low z , which would easily have been observed in surveys such as IRAS. And as shown in Fig. 1, even a significant fraction of galaxies at $z \lesssim 1$ would not match the measurements as the magnification boost is insufficient (again within the context of an underlying Schechter distribution).

4. DISCUSSION

We have considered the possibility that bright millimeter and submillimeter galaxy counts arise largely from a galaxy population at high redshifts that is lensed by intervening galaxy groups and clusters. Our model predictions match the counts from $500 \mu\text{m}$ to 2mm from the BLAST, SCUBA AzTEC and SPT surveys (see Fig. 2). We find that the high flux SPT number counts can be explained by highly magnified galaxies from this high- z population (once the known, low- z counterparts detected in IRAS are removed). This high- z galaxy pop-

ulation is described by a Schechter luminosity function with $L^* \sim 2.5 \times 10^{12} L_\odot$ and a median redshift of 3.0. Our model predictions fit the data for $500 \mu\text{m} < \lambda < 2 \text{mm}$ by varying S^* with wavelength within the range of typical submillimeter galaxy SEDs.

Our model has some simplifying assumptions, such as the fixed source redshift $z_s = 3$, and there are significant measurement and theoretical uncertainties. A complete analysis would require a more detailed treatment of the lensing and inclusion of the measured galaxy clustering. Nevertheless our results in Fig. 2 imply that current number counts do not require an additional galaxy population to explain the high flux measurements. Such a population has been invoked in theoretical models (Lagache et al. 2003) and suggested in the recent measurements of Vieira et al. (2009). Indeed, adding a significant fraction of a second population could cause the predictions to exceed the measured counts once magnification effects for the high- z population are included. It would also be difficult to explain how the scaling with wavelength of the high flux number counts is the same as the lower flux counts if they came from different galaxy populations. Since lensing does not depend on frequency, our model naturally follows this common scaling.

Current SZ surveys with sensitivities of $3 - 7 \times 10^{14} h^{-1} M_\odot$ cannot detect most halos that produce this lensing contribution. Conversely, looking for extremely bright objects in millimeter and submillimeter wavelengths can be an efficient way to search for high- z halos of masses $M \sim 10^{14} h^{-1} M_\odot$. The number of halos that can be found in this way is only a small fraction of all halos within a given mass range. Follow up observations with optical telescopes should be able to confirm cluster candidates up to $z \simeq 1$. These clusters and groups host Brightest Cluster Galaxies with luminosities $L \sim 10^{11} - 10^{12} L_\odot$, based on the low- z results of Johnston et al. (2007) from the Sloan Digital Sky Survey. Multi-band optical imaging with limiting magnitude of 23 – 24 (for the r band) would enable identification of these groups and clusters. Conversely, targeted observations of the high magnification regions of known strong lensing clusters could provide detections of faint submm galaxies (which would lie below the detection threshold without the magnification boost). Similar to the use of clusters as gravitational telescopes in optical imaging, this may also help resolve submm galaxies. Planned observations with AzTEC and the Large Millimeter Telescope have considered such an approach (David Hughes, private communication).

We thank David Hughes, Roxana Lupu, Joaquin Vieira, Kim Scott, Eric Switzer and various members of the ACT collaboration for useful discussions. We benefited from discussions on strong lensing effects with Matthias Bartelmann, Gary Bernstein, Neal Dalal and Ravi Sheth. We are very grateful to Stefan Hilbert for sharing his simulation results and Eric Switzer for providing the SPT data. This work was supported in part by an NSF-PIRE grant and AST-0607667.

REFERENCES

Aretxaga, I., et al. 2007, MNRAS, 379, 1571

- Austermann, J. E., et al. 2010, *MNRAS*, 401, 160
- Barger, A. J., Cowie, L. L., Sanders, D. B., Fulton, E., Taniguchi, Y., Sato, Y., Kawara, K., & Okuda, H. 1998, *Nature*, 394, 248
- Blain, A. W. 1996, *MNRAS*, 283, 1340
- Carlstrom, J. E., et al. 2009, ArXiv e-prints
- Chapman, S. C., Blain, A. W., Smail, I., & Ivison, R. J. 2005, *ApJ*, 622, 772
- Coppin, K., et al. 2006, *MNRAS*, 372, 1621
- Devlin, M. J., et al. 2009, *Nature*, 458, 737
- Dwek, E., et al. 1998, *ApJ*, 508, 106
- Eales, S., et al. 2009, *ApJ*, 707, 1779
- Fisher, K. B., Huchra, J. P., Strauss, M. A., Davis, M., Yahil, A., & Schlegel, D. 1995, *ApJS*, 100, 69
- Fixsen, D. J., Cheng, E. S., Gales, J. M., Mather, J. C., Shafer, R. A., & Wright, E. L. 1996, *ApJ*, 473, 576
- Fixsen, D. J., Dwek, E., Mather, J. C., Bennett, C. L., & Shafer, R. A. 1998, *ApJ*, 508, 123
- Fowler, J. W., et al. 2010, ArXiv e-prints
- Gonzalez, A. H., Clowe, D., Bradač, M., Zaritsky, D., Jones, C., & Markevitch, M. 2009, *ApJ*, 691, 525
- Greve, T. R., Ivison, R. J., Bertoldi, F., Stevens, J. A., Dunlop, J. S., Lutz, D., & Carilli, C. L. 2004, *MNRAS*, 354, 779
- Hilbert, S., White, S. D. M., Hartlap, J., & Schneider, P. 2007, *MNRAS*, 382, 121
- . 2008, *MNRAS*, 386, 1845
- Hincks, A. D., et al. 2008, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 7020, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series
- Hughes, D. H., et al. 1998, *Nature*, 394, 241
- Jain, B., & Lima, M. 2010, ArXiv e-prints
- Johnston, D. E., et al. 2007, ArXiv e-prints
- Kneib, J., van der Werf, P. P., Kraiberg Knudsen, K., Smail, I., Blain, A., Frayer, D., Barnard, V., & Ivison, R. 2004, *MNRAS*, 349, 1211
- Komatsu, E., et al. 2009, *ApJS*, 180, 330
- . 2010, ArXiv e-prints
- Lagache, G., Dole, H., & Puget, J. 2003, *MNRAS*, 338, 555
- Lagache, G., et al. 2004, *ApJS*, 154, 112
- Le Borgne, D., Elbaz, D., Ocvirk, P., & Pichon, C. 2009, *A&A*, 504, 727
- Lima, M., Jain, B., & Devlin, M. 2009, ArXiv e-prints
- Lima, M., et al. 2010, in prep.
- Moshir, M., Kopman, G., & Conrow, T. A. O. 1992, *IRAS Faint Source Survey, Explanatory supplement version 2*, ed. Moshir, M., Kopman, G., & Conrow, T. A. O.
- Negrello, M., Perrotta, F., González-Nuevo, J., Silva, L., de Zotti, G., Granato, G. L., Baccigalupi, C., & Danese, L. 2007, *MNRAS*, 377, 1557
- Oliver, S. J., et al. 1996, *MNRAS*, 280, 673
- Paciga, G., Scott, D., & Chapin, E. L. 2008, ArXiv e-prints
- Pearson, C., & Khan, S. A. 2009, *MNRAS*, 399, L11
- Pérez-González, P. G., et al. 2005, *ApJ*, 630, 82
- Perrotta, F., Baccigalupi, C., Bartelmann, M., De Zotti, G., & Granato, G. L. 2002, *MNRAS*, 329, 445
- Pope, A., et al. 2006, *MNRAS*, 370, 1185
- Rex, M., et al. 2009, *ApJ*, 703, 348
- Schechter, P. 1976, *ApJ*, 203, 297
- Smail, I., Ivison, R. J., & Blain, A. W. 1997, *ApJ*, 490, L5+
- Smail, I., Ivison, R. J., Blain, A. W., & Kneib, J. 2002, *MNRAS*, 331, 495
- Staniszewski, Z., et al. 2009, *ApJ*, 701, 32
- Vanderlinde, K., et al. 2010, ArXiv e-prints
- Vieira, J. D., et al. 2009, ArXiv e-prints
- Wilson, G. W., et al. 2008, *MNRAS*, 390, 1061